

素, 考虑到这些因素多与高血压有密切联系, 因而推测它们是通过影响血压而间接地促使了缺血性脑卒中的发生。

A Study on Risk Factors of Ischemic Stroke Qiu Chengxuan, et al., Department of Epidemiology, Tianjin Medical College, Tianjin

In order to investigate risk factors of ischemic stroke, a matched case-control study with two kinds of controls (community controls and hospital controls) was conducted. The multiple conditional logistic regression analysis showed that hypertension, transient ischemic attacks, alcohol consumption and habitual snoring were independent risk factors of ischemic stroke. Physical activity at leisure time was an important protective factor. No independent relationship were found between cigarette smoking, history of mellitus diabetes or heart diseases, stroke history of father or mother and ischemic stroke.

Key words Case-control study Conditional logistic regression Ischemic stroke

参 考 文 献

1. Camargo Jr. CA. Moderate alcohol consumption

and stroke: the epidemiology evidence. Stroke 1989; 20: 1611.

2. 各类脑血管疾病诊断要点(1986年中华医学会第二次脑血管病学术会议第三次修订). 中华神经精神科杂志 1988; 21: 60.

3. Doll R and Hill AB. Smoking and carcinoma of the lung preliminary report. Br Med J 1950; 2: 793.

4. Silverman DG, et al. Artificial sweeteners and lower urinary tract cancer: Hospital vs population controls. Am J Epidemiol 1983; 117: 326.

5. Report of the WHO task force on stroke and other cerebrovascular diseases (WHO). Stroke 1989; Recommendations on stroke preventions, diagnosis, and therapy. Stroke 1989; 20: 1407.

6. Davis PH, et al. Risk factors for ischemic stroke: a prospective study in Rochester, Minnesota. Ann Neurol 1987; 22: 319.

7. Partinen M and Palomaki H. Snoring and cerebral infarction. Lancet 1985; ii: 1325.

8. Herman B, et al. Multivariate logistic analysis of risk factors for stroke in Tilburg, the Netherlands. Am J Epidemiol 1983; 118: 514.

9. Wolf PA, et al. Cigarette smoking as a risk factor for stroke: the Framingham Study. JAMA 1988; 259: 1025.

(1990年10月19日收稿, 1992年8月4日修回)

**葡萄糖胰岛素氯化钾三联液治疗肝病
蛋白质代谢异常的初步观察**

辽宁省辽阳市传染病院* 洪 明 朴盛琴 唐素梅 张晓东

应用葡萄糖胰岛素氯化钾三联液治疗急、慢性肝病蛋白质代谢异常, 通过一个疗程的治疗观察效果显著, 血浆总蛋白治疗前为 $63 \pm 5\text{g/L}$, 治疗后为 $72.9 \pm 6\text{g/L}$, 疗后较疗前明显提高 ($P < 0.05$); 血清白蛋白治疗前为 $34.5 \pm 10\text{g/L}$, 治疗后为 $46.3 \pm 5\text{g/L}$, 治疗后提升明显 ($P < 0.01$); 球蛋白治疗前为 $34 \pm 10\text{g/L}$,

治疗后为 $25.8 \pm 4\text{g/L}$, 疗后较疗前也有明显降低 ($P < 0.05$)。结果表明该三联液在调整血浆蛋白、纠正白球蛋白比例倒置方面有着明显效果, 而且不受条件限制, 便于基层应用, 并且价格便宜。这对解决当前肝病用药价格昂贵等情况也是一个良好的开端。

* 邮政编码 111000