

- [9] Guangdong Provincial Health Commission. The first imported case of COVID-19 by National Health Commission confirmed in Guangdong province[EB/OL]. (2020-01-20) [2020-11-20]. http://wsjkw.gd.gov.cn/zwy_yqxx/content/post_2876057.html.
- [10] 国家卫生健康委员会.国家卫生健康委员会公告[EB/OL]. (2020-01-20) [2020-11-20]. <http://www.nhc.gov.cn/jkj/s7916/202001/44a3b8245e8049d2837a4f27529cd386.shtml>. National Health Commission. Announcement of the National Health Commission [EB/OL]. (2020-01-20) [2020-11-20]. <http://www.nhc.gov.cn/jkj/s7916/202001/44a3b8245e8049d2837a4f27529cd386.shtml>.
- [11] 国家卫生健康委员会办公厅.新型冠状病毒肺炎诊疗方案(试行第八版)[J].中国医药, 2020, 15(10):1494-1499. DOI:10.3760/j.issn.1673-4777.2020.10.002.
- [12] General Office of National Health Commission, Office of National Administration of Traditional Chinese Medicine. Protocol of diagnosis and treatment for COVID-19(trial version 8) [J]. China Med, 2020, 15(10):1494-1499. DOI:10.3760/j.issn.1673-4777.2020.10.002.
- [13] 国家卫生健康委员会办公厅.新型冠状病毒肺炎防控方案(第七版)的通知[EB/OL]. (2020-09-15) [2021-01-27]. <http://www.nhc.gov.cn/jkj/s3577/202009/318683cbfaee4191aee29cd774b19d8.shtml>.
- [14] General Office of National Health Commission. Protocol of prevention and control for COVID-19(version 7) [EB/OL]. (2020-09-15) [2021-01-27]. <http://www.nhc.gov.cn/jkj/s3577/202009/318683cbfaee4191aee29cd774b19d8.shtml>.
- [15] Cheng VCC, Wong SC, Chuang VWM, et al. The role of community-wide wearing of face mask for control of coronavirus disease 2019 (COVID-19) epidemic due to SARS-CoV-2[J]. J Infect, 2020, 81(1): 107-114. DOI: 10.1016/j.jinf.2020.04.024.
- [16] Zhong BL, Luo W, Li HM, et al. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey[J]. Int J Biol Sci, 2020, 16(10):1745-1752. DOI:10.7150/ijbs.45221.
- [17] Ngewendo A, Nkengazong L, Ambe LA, et al. Knowledge, attitudes, practices of/towards COVID 19 preventive measures and symptoms: A cross-sectional study during the exponential rise of the outbreak in Cameroon[J]. PLoS Negl Trop Dis, 2020, 14(9): e0008700. DOI: 10.1371/journal.pntd.0008700.
- [18] Wake AD. Knowledge, attitude, practice, and associated factors regarding the novel coronavirus disease 2019 (COVID-19) pandemic[J]. Infect Drug Resist, 2020, 13: 3817-3832. DOI:10.2147/IDR.S275689.
- [19] Morawska L, Tang JW, Bahnfleth W, et al. How can airborne transmission of COVID-19 indoors be minimised? [J]. Environ Int, 2020, 142: 105832. DOI: 10.1016/j.envint.2020.105832.
- [20] Xia XY, Wu J, Liu HL, et al. Epidemiological and initial clinical characteristics of patients with family aggregation of COVID-19[J]. J Clin Virol, 2020, 127: 104360. DOI: 10.1016/j.jcv.2020.104360.
- [21] Ghinai I, Woods S, Ritger KA, et al. Community transmission of SARS-CoV-2 at two family gatherings - Chicago, Illinois, February-March 2020[J]. MMWR Morb Mortal Wkly Rep, 2020, 69(15):446-450. DOI:10.15585/mmwr.mm6915e1.
- [22] Cheng HY, Jian SW, Liu DP, et al. Contact tracing assessment of COVID-19 transmission dynamics in Taiwan and risk at different exposure periods before and after symptom onset[J]. JAMA Intern Med, 2020, 180(9): 1156-1163. DOI:10.1001/jamainternmed.2020.2020.
- [23] Li W, Zhang B, Lu JH, et al. Characteristics of household transmission of COVID-19[J]. Clin Infect Dis, 2020, 71(8): 1943-1946. DOI:10.1093/cid/ciaa450.
- [24] Grijalva CG, Rolfes MA, Zhu YW, et al. Transmission of SARS-CoV-2 infections in households-Tennessee and Wisconsin, April-September 2020[J]. MMWR Morb Mortal Wkly Rep, 2020, 69(44):1631-1634. DOI:10.15585/mmwr.mm6944e1.
- [25] 张文佳,廖璞.1例新型冠状病毒肺炎多次核酸检测异同结果的原因分析[J].重庆医学, 2020, 49(17):2834-2836. DOI:10.3969/j.issn.1671-8348.2020.17.013.
- [26] Zhang WJ, Liao P. Analysis of the reasons for similarities and differences of multiple times of nucleic acid testing for one case of COVID-19[J]. Chongqing Med, 2020, 49 (17): 2834-2836. DOI: 10.3969/j. issn. 1671-8348. 2020. 17.013.
- [27] 关明,王文静,王惠英.新型冠状病毒无症状感染者的实验室检测及防控风险评估[J].中华预防医学杂志, 2020, 54 (12):1310-1314. DOI:10.3760/cma.j.cn112150-20200610-00861.
- [28] Guan M, Wang WJ, Wang HY. Laboratory testing, prevention and control of asymptomatic carriers of coronavirus disease 2019 [J]. Chin J Prev Med, 2020, 54 (12):1310-1314. DOI: 10.3760/cma.j.cn112150-20200610-00861.
- [29] 李振昊,高小玲,杨小姐,等.新型冠状病毒核酸检测分析[J].检验医学与临床, 2020, 17(10): 1313-1315. DOI: 10.3969/j.issn.1672-9455.2020.10.001.
- [30] Li ZH, Gao XL, Yang XJ, et al. Detection and analysis of SARS-CoV-2 nucleic acid[J]. Lab Med Clinic, 2020, 17(10): 1313-1315. DOI:10.3969/j.issn.1672-9455.2020.10.001.
- [31] 郑甲兰.PCR检测中的假阳性和假阴性问题分析[J].中国医药指南, 2012, 10(22):390-391. DOI:10.3969/j.issn.1671-8194.2012.22.296.
- [32] Zheng JL. Analysis of false positive and false negative in PCR detection[J]. Guide China Med, 2012, 10(22): 390-391. DOI:10.3969/j.issn.1671-8194.2012.22.296.
- [33] Hao W, Ma B, Li ZH, et al. Binding of the SARS-CoV-2 spike protein to glycans[J]. Sci Bull, 2021. DOI: 10.1016/j.scib.2021.01.010.
- [34] Rajawat J, Chandra A. Role of Poly (ADP-ribose) polymerase (PARP1) in viral infection and its implication in SARS-CoV-2 pathogenesis[J]. Curr Drug Targets, 2021. DOI:10.2174/1389450122666210120142746.
- [35] Corley MJ, Pang APS, Dody K, et al. Genome-wide DNA methylation profiling of peripheral blood reveals an epigenetic signature associated with severe COVID-19[J]. J Leukoc Biol, 2021. DOI:10.1002/JLB.5HI0720-466R.
- [36] Combes AJ, Courau T, Kuhn NF, et al. Global absence and targeting of protective immune states in severe COVID-19[J]. Nature, 2021. DOI: 10.1038/s41586-021-03234-7.
- [37] Quilty BJ, Clifford S, Hellewell J, et al. Quarantine and testing strategies in contact tracing for SARS-CoV-2: a modelling study[J]. Lancet Public Health, 2021. DOI: 10.1016/S2468-2667(20)30308-X.
- [38] Spencer KD, Chung CL, Stargel A, et al. COVID-19 case investigation and contact tracing efforts from health departments-United States, June 25-July 24, 2020[J]. MMWR Morb Mortal Wkly Rep, 2021, 70(3):83-87. DOI: 10.15585/mmwr.mm7003a3.
- [39] Liu HZ, Ye CC, Wang YP, et al. The effectiveness of active surveillance measures for COVID-19 cases in Pudong New Area Shanghai, China, 2020[J]. J Med Virol, 2021. DOI: 10.1002/jmv.26805.