











能导致未报告的疑似麻疹暴发起数被高估。

在全球致力于消除麻疹的现阶段,进一步做好麻疹监测和暴发疫情调查处置更为重要。WHO 于 2021 年发布麻疹暴发疫情处置应对策略指南,在加强基础免疫及提升麻疹疫苗第二剂次接种率、减少不同地区免疫水平差异防止疫情暴发、加强监测和及时有效地处置疫情等方面提出了建议<sup>[21]</sup>。结合我国现状,进一步做好麻疹暴发疫情深入调查、及时报告并落实处置措施,仍是下一步消除麻疹工作的重点。

利益冲突 所有作者均声明不存在利益冲突

### 参 考 文 献

- [1] 马超,贾海梅,苏琪茹,等.中国 2009-2015 年麻疹暴发疫情报告与调查处置情况分析[J].中国疫苗和免疫,2016,22(5):481-486,492.  
Ma C, Jia HM, Su QR, et al. Surveillance and response to measles outbreaks in China[J]. Chin J Vacc Immun, 2016, 22(5):481-486, 492.
- [2] 马超,苏琪茹,郝利新,等.中国 2017-2018 年麻疹流行病学特征[J].中国疫苗和免疫,2020,26(1):5-8,41.  
Ma C, Su QR, Hao LX, et al. Epidemiology of measles in China, 2017-2018[J]. Chin J Vacc Immun, 2020, 26(1): 5-8, 41.
- [3] Patel MK, Goodson JL, Alexander JP, et al. Progress toward regional measles elimination-worldwide, 2000-2019[J]. MMWR Morb Mortal Wkly Rep, 2020, 69(45):1700-1705. DOI:10.15585/mmwr.mm6945a6.
- [4] WHO. Measles elimination field guide[M]. Manila: WHO Regional Office for the Western Pacific, 1993.
- [5] 马超,郝利新,温宁,等.中国 2019 年麻疹流行病学特征[J].中国疫苗和免疫,2020,26(5):493-497.  
Ma C, Hao LX, Wen N, et al. Epidemiology of measles in China, 2019[J]. Chin J Vacc Immun, 2020, 26(5):493-497.
- [6] 疾病预防控制中心.2020 年全国法定传染病疫情概况[EB/OL].(2021-03-12) [2021-05-01]. <http://www.nhc.gov.cn/jkj/s3578/202103/f1a448b7df7d4760976fea6d55834966.shtml>.  
Bureau of Disease Control and Prevention. Overview of national epidemic situation of legal infectious diseases in 2020[EB/OL]. (2021-03-12) [2021-05-01]. <http://www.nhc.gov.cn/jkj/s3578/202103/f1a448b7df7d4760976fea6d55834966.shtml>.
- [7] 卫生部.卫生部办公厅关于印发《全国麻疹监测方案》的通知[EB/OL].(2009-02-02) [2021-05-01]. <http://www.nhc.gov.cn/zwgk/wtwj/201304/f9956cbc95594311902cf554e0aee4b7.shtml>.  
Ministry of Health. Notice of the General Office of the Ministry of Health on printing and distributing the National Measles Monitoring Program[EB/OL]. (2009-02-02) [2021-05-01]. <http://www.nhc.gov.cn/zwgk/wtwj/201304/f9956cbc95594311902cf554e0aee4b7.shtml>.
- [8] Ma C, Rodewald L, Hao LX, et al. Progress toward measles elimination-China, January 2013-June 2019[J]. MMWR Morb Mortal Wkly Rep, 2019, 68(48):1112-1116. DOI:10.15585/mmwr.mm6848a2.
- [9] Guerra FM, Bolotin S, Lim G, et al. The basic reproduction number (R0) of measles: a systematic review[J]. Lancet Infect Dis, 2017, 17(12):e420-428. DOI:10.1016/S1473-3099(17)30307-9.
- [10] Hao LX, Glasser JW, Su QR, et al. Evaluating vaccination policies to accelerate measles elimination in China: a meta-population modelling study[J]. Int J Epidemiol, 2019, 48(4):1240-1251. DOI:10.1093/ije/dyz058.
- [11] Misin A, Antonello RM, Di Bella S, et al. Measles: an overview of a re-emerging disease in children and immunocompromised patients[J]. Microorganisms, 2020, 8(2):276. DOI:10.3390/microorganisms8020276.
- [12] Harapan H, Itoh N, Yufika A, et al. Coronavirus disease 2019 (COVID-19): A literature review[J]. J Infect Public Health, 2020, 13(5):667-673. DOI:10.1016/j.jiph.2020.03.019.
- [13] 马超,苏琪茹,温宁,等.中国 2015-2016 年麻疹监测系统运转情况分析[J].中国疫苗和免疫,2018,24(2):141-145.  
Ma C, Su QR, Wen N, et al. Evaluation of the measles surveillance system performance in China, 2015-2016 [J]. Chin J Vacc Immun, 2018, 24(2):141-145.
- [14] 沙吾拉西·热加甫,玛合木提江·库尔班,谢娜,等.2012-2018 年新疆维吾尔自治区麻疹流行病学特征分析[J].疾病监测,2019,34(12):1109-1112. DOI:10.3784/j.issn.1003-9961.2019.12.016.  
Rejiafu S, Kuerban M, Xie N, et al. Epidemiological characteristics of measles in Xinjiang, 2012-2018[J]. Dis Surveill, 2019, 34(12):1109-1112. DOI:10.3784/j.issn.1003-9961.2019.12.016.
- [15] Remington PL, Hall WN, Davis IH, et al. Airborne transmission of measles in a physician's office[J]. JAMA, 1985, 253(11):1574-1577. DOI:10.1001/jama.1985.03350350068022.
- [16] Hope K, Boyd R, Conaty S, et al. Measles transmission in health care waiting rooms: implications for public health response[J]. Western Pac Surveill Response J, 2012, 3(4):33-38. DOI:10.5365/WPSAR.2012.3.3.009.
- [17] Moss WJ, Griffin DE. Measles[J]. Lancet, 2012, 379(9811):153-164. DOI:10.1016/S0140-6736(10)62352-5.
- [18] Fine PEM, Clarkson JA. Measles in England and Wales- I : An analysis of factors underlying seasonal patterns[J]. Int J Epidemiol, 1982, 11(1):5-14. DOI:10.1093/ije/11.1.5.
- [19] Chen M, Zhang Y, Huang F, et al. Endemic and imported measles virus-associated outbreaks among adults, Beijing, China, 2013[J]. Emerg Infect Dis, 2015, 21(3):477-479. DOI:10.3201/eid2103.140646.
- [20] 王淑蕾,李崇山,王慧玲,等.中国大陆首次发现输入性 B3 基因型麻疹病毒病例[J].病毒学报,2014,30(5):535-540. DOI:10.13242/j.cnki.bingduxuebao.002544.  
Wang SL, Li CS, Wang H et al. Imported B3 genotype measles viruses were isolated from measles cases in the Chinese mainland[J]. Chin J Virol, 2014, 30(5):535-540. DOI:10.13242/j.cnki.bingduxuebao.002544.
- [21] World Health Organization. Measles outbreaks strategic response plan: 2021-2023: measles outbreak prevention, preparedness, response and recovery[EB/OL]. (2020-10-07) [2021-06-25] <https://apps.who.int/iris/handle/10665/340657>.